

MDR-LIKE ABC TRANSPORTER GENE FROM PLANTS

This application is a continuation in part of International Application No. PCT/US99/22363, filed September 24, 1999, which claims priority under 35 U.S.C. §120 to U.S. Provisional Application 60/101,814, the entireties of both of which are incorporated by reference herein.

Pursuant to 35 U.S.C. §202(c), it is acknowledged that the U.S. Government has certain rights in the invention described herein, which was made in part with funds from the National Science Foundation, Grant No. IBN-9416016.

FIELD OF THE INVENTION

This invention relates to the field of stress resistance in plants. In particular, the invention provides a novel gene in plants, which encodes an MDR-like ABC transporter involved in detoxification of certain xenobiotics to protect plants from their detrimental effects.

BACKGROUND OF THE INVENTION

Several applications are referenced in this background section, which are hereby incorporated by reference.

The following references are cited in the background section of this application, which are hereby incorporated by reference.

the stress caused by environmental pollutants in the soil, water and atmosphere. Such pollutants include herbicides, pesticides and related agronomic products, as well as organic and inorganic waste material from industry and other sources.

5 Other toxic agents that threaten the survival of plants include various toxins produced by epiphytic or soilborne microorganisms, such as fungi and bacteria.

In order to survive in toxic environments, plants must have mechanisms to detoxify xenobiotics, heavy metals
10 and other toxic compounds. This generally involves modification of the toxic compound and subsequent excretion into the vacuole or apoplastic space. Recently, certain ATP-binding cassette (ABC) transporters have been identified in plants, which appear to be involved in the detoxification
15 process.

The ABC transporter family is very large, with representatives existing in many different classes of organisms. Two well studied groups of ABC transporters, encoded by *mdr* and *mrp* genes, respectively, are associated
20 with the multidrug resistance phenomenon observed in mammalian tumor cells. The *mdr* genes encode a family of P-glycoproteins that mediate the energy-dependent efflux of certain lipophilic drugs from cells. The *mrp* genes encode a family of transporters that mediate the extrusion of a
25 variety of hydrophobic compounds after their conjugation with

glutathione. *mdr* and *mrp* genes have been identified

conjugate transporter encoded by the *mrp* homolog is located in the vacuolar membrane and is responsible for sequestration of xenobiotics in the central vacuole (Tommasini et al., FEBS Lett. 411: 306-310, 1997; Id et al., Plant Physiol. 107:

5 1257-1264, 1993). An *mdr*-like gene (*atpgp1*) has also been identified in *A. thaliana*, which encodes a putative P-glycoprotein homolog. The *atpgp1* gene was found to share significant sequence homology and structural organization with human *mdr* genes, and was expressed with particular
10 abundance in inflorescence axes (Dudler & Hertig, J. Biol. Chem. 267: 4881-4888, 1992). Other MDR homologs have been found in potato (Wang et al., Plant Mol. Biol. 31: 683, 1996) and barley (Davies et al., Gene 199: 195, 1997).

The aforementioned *mrp* and *mdr* plant homologs were
15 identified as a result of an effort to understand the molecular basis for development in plants of cross-resistance to herbicides of unrelated classes. However, these transporters are likely to serve the more general role in plants of sequestering, secreting, or otherwise detoxifying
20 various xenobiotics and endogenous xenobiotics. Accordingly, it will continue to be an area in the art of plant genetic engineering to which tolerance to identify and characterize other members of this class of transporters in plants.

25 SUMMARY OF THE INVENTION

The present invention provides a method for identifying a class of transporters

inducible by NPPB and binary NPA.

According to one aspect of the invention, a nucleic acid isolated from a plant is provided, which encodes a p-glycoprotein that is inducible by exposure of the plant to NPPB or NPA. The isolated nucleic acid is preferentially expressed in plant roots upon exposure of the plant to NPPB. In a preferred embodiment, the plant from which the nucleic acid is isolated is selected from the group consisting of *Brassica napus* and *Arabidopsis thaliana* and is 3850-4150 nucleotides in length. In a more preferred embodiment, the nucleic acid has the restriction sites shown in Figure 4 for at least three restriction enzymes. In particularly preferred embodiments, the nucleic acid molecule encodes a polypeptide having SEQ ID NO:2. In an exemplary embodiment, the nucleic acid is a cDNA comprising the coding region of SEQ ID NO:1 or SEQ ID NO:10.

According to another aspect of the invention is an expression cassette that comprises a pLPAC gene operably linked to a promoter, and in a more preferred embodiment the pLPAC gene is from *Arabidopsis*. In preferred embodiments, the expression cassette comprises the cauliflower mosaic virus 35S promoter, and part or all of SEQ ID NO:1 or SEQ ID NO:10. Further included in this aspect is a vector comprising the expression cassette and a method for producing transgenic plants with the expression cassette and vector.

According to a further aspect of the invention, a transgenic plant is provided which contains the expression cassette.

the transmission ent.

According to another aspect of the invention, an isolated nucleic acid molecule is provided, which has a sequence selected from the group consisting of: a) SEQ ID NO:1 and SEQ ID NO:10; b) a nucleic acid sequence that is at least about 60% homologous to the coding regions of SEQ ID NO:1 or SEQ ID NO:10; c) a sequence hybridizing with SEQ ID NO:1 or SEQ ID NO:10 at moderate stringency; d) a sequence encoding part or all of a polypeptide having SEQ ID NO:2; e) a sequence encoding an amino acid sequence that is at least about 70% identical to SEQ ID NO:2; f) a sequence encoding an amino acid sequence that is at least about 80% similar to SEQ ID NO:1; g) a sequence encoding an amino acid sequence that is at least about 40% similar to residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2; and h) a sequence hybridizing at moderate stringency to a sequence encoding residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2. A polypeptide produced by expression of the above listed sequences is also provided.

According to another aspect of the invention, an isolated plant polypeptide, which is inducible upon exposure to the herbicide NPPB, is provided. The polypeptide preferably is isolated from a plant in which it is found resistant to the herbicide 63. The polypeptide is preferably synthesized in plants upon the exposure to the NPPB. The plant from which it is preferably from *Brassica napus* or

1-76, 613-669 or 1144-1161 of SEQ ID NO:2; and d) an amino acid sequence encoded by a nucleic acid sequence hybridizing at moderate stringency to an amino acid sequence encoding residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2.

5 According to other aspects of the invention, antibodies immunologically specific for the polypeptides of the invention are provided, that immunologically specific to any of the polypeptides, of polypeptide encoded by the nucleic acids of the invention. In a preferred embodiment,
10 the antibody is immunospecific to residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2.

According to another aspect of the invention, a plant polypeptide gene promoter, which is inducible by NPPB, is also provided. In a preferred embodiment, the
15 promoter is part or all of residues 1-3429 of SEQ ID NO:10. According to another aspect of the invention, plants that have reduced levels of pLPAC protein are provided. In a preferred embodiment, these plants have mutations in the pLPAC gene, and in a particularly preferred embodiment, the
20 pLPAC gene contains the insertion of a T-DNA. Also provided is a method for selecting plants with mutations in pLPAC using SEQ ID NOS:11-14 as PCR primers.

These and other features and advantages of the
25 present invention will be described in greater detail in the

BRIEF DESCRIPTION OF DRAWINGS

Figure 1. Nucleotide sequence (nt) : ATTA

related mammalian and plant genes. The lineup shows the ATPAC deduced amino acid sequence (SEQ ID NO:2) compared with (1) hmdr1 (SEQ ID NO:3); (2) hmdr1 (SEQ ID NO: 4); (3) hmdr3 (SEQ ID NO:5); (4) hmdr2 (SEQ ID NO:6); (5) atpgp1 (SEQ ID NO:7); and (6) atpgp1 (SEQ ID NO:8). A consensus sequence (SEQ ID NO:9) is also shown.

Figure 2. Graph depicting the effect of rhodamine 6G on the growth rate of cells transformed with and expressing ATPAC as compared with control cells not containing ATPAC.

Figure 3. Restriction map of genomic clone of ATPAC, SEQ ID NO:10.

Figure 4. Restriction map of cDNA clone of ATPAC, SEQ ID NO:11.

DETAILED DESCRIPTION OF THE INVENTION

I. Definitions

Various terms relating to the biological molecules of the present invention are used hereinabove and also through the description and claims.

With respect to nucleic acids of the invention, the term "nucleic acid" is sometimes used. This term, when applied to DNA, refers to a DNA molecule that is separate from sequences with which it is immediately contiguous (in opposite directions) in the naturally

occurring DNA. The term "genomic DNA" refers to a prokaryotic or eukaryotic genomic DNA molecule.

comprise a DNA molecule.

With respect to RNA molecules of the invention the term "isolated nucleic acid" primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

Nucleic acid sequences and amino acid sequences can be compared using computer programs that align the similar sequences of the nucleic or amino acids thus define the differences. For purposes of this invention, the DNASTar program (DNASTar, Inc., Madison, Wisconsin) and the default parameters used by that program are the parameters intended to be used herein to compare sequence identity and similarity. Alternately, the Blastn and Blastp 2.0 programs provided by the National Center for Biotechnology Information (at <http://www.ncbi.nlm.nih.gov/blast/>; Altschul et al., 1990, J. Mol. Biol. 214:403-416) using a gapped alignment with default parameters, may be used to determine the level of identity and similarity between nucleic acid sequences and amino acid sequences.

The term "substantially the same" refers to nucleic acid or amino acid sequences having sequence variation that

under the term "substantially the same" is

sequence, governing expression, and refers primarily to degenerate codons encoding the same amino acid, or alternate codons encoding conservative substitute amino acids in the encoded polypeptide. With reference to amino acid sequences, the term "substantially the same" refers generally to conservative substitutions and/or variations in regions of the polypeptide involved in determination of structure or function.

The terms "percent identical" and "percent similar" are also used herein in comparisons among amino acid and nucleic acid sequences. When referring to amino acid sequences, "percent identical" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical amino acids in the compared amino acid sequence by a sequence analysis program. "Percent similar" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical or conservative amino acids. Conserved amino acids are those which differ in structure but are similar in physical properties such that they could be substituted for another would not appreciably change the tertiary structure of the resulting protein. Conservative substitutions are defined in Taylor (1986, J. Theor. Biol. 119:205). When referring to nucleic acid molecules, "percent identical" refers to the percent of the nucleotides of the subject nucleic acid sequence that

expression of an isolated nucleic acid molecule of the invention. Alternatively, this term may refer to a protein which has been sufficiently separated from other proteins with which it would naturally be associated, so as to exist in "substantially pure" form.

The term "substantially pure" refers to a preparation comprising at least 50-60% by weight the compound of interest (e.g., nucleic acid, oligonucleotide, protein, etc.). More preferably, the preparation comprises at least 75% by weight, and most preferably 90-99% by weight, the compound of interest. Purity is measured by methods appropriate for the compound of interest (e.g., chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

With respect to antibodies of the invention, the term "immunologically specific" refers to antibodies that bind to one or more epitopes of a protein of interest, but which do not substantially recognize and bind other molecules in a sample containing a mixed population of antigenic biological molecules.

With respect to nucleotides, the term "specific hybridizing" refers to the association between two single-stranded nucleotide molecules of sufficiently complementary sequence to permit such hybridization under pre-determined conditions generally used in the art.

It is to be understood that the term "nucleic acid molecule" as used herein includes both DNA and RNA molecules and that the term "nucleotide" as used herein includes both deoxyribonucleotides and ribonucleotides.

the oligonucleotide with single-stranded nucleic acids of non-complementary sequence.

The term "expression cassette", as used herein, comprises at least one regulatory region operably linked to a coding sequence. The coding sequence may be in the sense or antisense orientation with respect to the 5' regulatory region.

The term "promoter region" refers to the 5' regulatory region of a gene.

The term "reporter gene" refers to genetic sequences which may be operably linked to a promoter region forming a transgene, such that expression of the reporter gene coding region is regulated by the promoter and expression of the transgene is readily assayed.

The term "selectable marker gene" refers to a gene product that when expressed confers a selectable phenotype, such as antibiotic resistance, on a transformed cell or plant.

The term "operably linked" means that the regulatory sequence necessary for expression of the coding sequence is placed in the DNA molecule in the appropriate position relative to the coding sequence so as to effect expression of the coding sequence. This same definition is sometimes applied to the arrangement of coding sequences and transcription control elements (e.g. promoters, enhancers,

plant growth regulators, etc.) administered to plants in a

Agrobacterium T-DNA mediated transformation and transformation using the biolistic process are also contemplated to be within the scope of the present invention. The transforming DNA may be prepared according to standard protocols such as those set forth in "Current Protocols in Molecular Biology", eds. Frederick M. Ausubel et al., John Wiley & Sons, 2000.

The term "xenobiotic" refers to foreign chemicals or agents not produced or naturally found in the organism. The term is commonly used in reference to toxic or otherwise detrimental foreign chemicals, such as organic pollutants or heavy metals.

II. Description of *pIPAC* and its Encoded Polypeptide

In accordance with the present invention, a nucleic acid encoding a novel ATP-binding-cassette (ABC) transporter has been isolated and cloned from plants. This novel ABC transporter is encoded by *atpabc* and binds NPA. The nucleic acid is referred to herein as *pIPAC*.

Sequence of the *pIPAC* from *Arabidopsis thaliana*, an example of *pIPAC* of the invention, is described in detail herein. Its nucleotide sequence is set forth in Example 1 of SEQ. ID. NO. 1. This nucleic acid molecule is referred to as "1.1". It is 46% identical and 51% similar

to the sequence of the *atpabc* gene from *Arabidopsis thaliana* (GenBank accession number U00096). *atpabc* is a member of the ABC transporter family. *atpabc* protein is

A part of the cDNA of a *plPAC* of the invention was originally isolated from *Brassica napus* via differential expression between plants grown in the presence or absence of the chloride channel blocker, 5-nitro-2-(3-phenylpropyl)pyrimidin-4-carboxylic acid (NPPB). A 0.5 kb gene fragment was identified, which had been up-regulated in response to NPPB treatment. This cDNA fragment was used to screen an *Arabidopsis* cDNA library, from which the complete *ATPAC* cDNA was isolated. The isolation and characterization of *ATPAC* is described in Example 1.

A genomic clone of *ATPAC* (SEQ ID NO:10) has also been isolated from a bacterial artificial chromosome (BAC) library of the *Arabidopsis* genome (BAC clone IGF F3J22, obtained from the *Arabidopsis* stock center, Ohio State University). A 1.8 kb fragment containing part of *ATPAC* and additional 5' regulatory sequences was subcloned into a plasmid vector (pBluescript). A restriction map of *ATPAC* is found in Fig. 3. The corresponding cDNA clone of *ATPAC* is found in SEQ ID NO:11 and its restriction map is Fig. 4.

In *Arabidopsis*, *plPAC* of the present invention is expressed in roots, leaves, flowers, and shoot meristem. Expression is induced in the hypocotyl of etiolated seedlings, but not in light-grown seedlings. Expression of *ATPAC* is also relatively high in cotyledons, meristem, roots and the first true leaves of the seedling.

When the *ATPAC* gene is expressed in *Arabidopsis*, the potential expression in

residues 1-429 of SEQ ID NO:10.

Although the ATHAS cDNA clone from *Arabidopsis thaliana* is described and exemplified herein, this invention is intended to encompass nucleic acid sequences and proteins from other plant species that are sufficiently similar to be used instead of ATHAS nucleic acid and proteins for the purposes described below. These include, but are not limited to, allelic variants and natural mutants of SEQ ID NO:1, which are likely to be found in different species of plants or varieties of *Arabidopsis*.

Because such variants are expected to possess certain differences in nucleotide and amino acid sequence, this invention provides an isolated *pIPAC* nucleic acid molecule having at least about 60% (preferably 70% and more preferably over 80% sequence homology in the coding regions with the nucleotide sequence set forth as SEQ ID NO:1 or SEQ ID NO:10 and, most preferably, specifically comprising the coding region of SEQ ID NO:1). Also provided are nucleic acids that encode a polypeptide that is at least about 40% (preferably 50% and most preferably 60%) similar to residues 1-76, 81-113 or 114-116 of SEQ ID NO:2. Also provided are nucleic acids that are homologous to the nucleic acids of SEQ ID NO:1, SEQ ID NO:10, or nucleic acids encoding the regions of residues 1-76, 81-113 or 114-116 of SEQ ID NO:2, preferably with a high degree of stringency (more preferably, high

stringency) and which are at least 50% and most preferably 70% or more similar to the sequence of SEQ ID NO:1 or SEQ ID NO:10.

preferred embodiment, the nucleic acids have a restriction digest map identical to those shown in Fig. 3 for enzymes XhoI, XbaI and SalI (preferably additionally SacI, PacI and BsaI, and most preferably additionally AclI, BanI and SnaBI).

5 In another preferred embodiment, the nucleic acids have a restriction digest map identical to those shown in Fig. 4 for enzymes XbaI, TatI and NciI (preferably additionally DraI, PvuI and BclI, and most preferably additionally AclI, AclI and TliI). The nucleic acids of the invention are at least 20 nucleic acids in length (preferably
10 at least 50 nucleic acids and most preferably at least 100 nucleic acids).

In accordance with the invention, novel pLPAC genes from two plant species, *Brassica napus* and *Arabidopsis*
15 *thaliana*, are presented. This constitutes the first description of this unique p-glycoprotein in plants. Indeed, the closest known protein sequence, also from *Arabidopsis*, is only 65% identical, suggesting that the ATPAC gene is novel and is expected to have novel properties. The isolation of
20 two pLPAC genes from different species enables the isolation of further pLPAC genes from other plant species. Isolated nucleic acid sequences of pLPAC genes from any plant species are considered part of the instant invention. In particular, the nucleic acid sequences of pLPAC genes can be isolated using
25 sequences of ATPAC to distinguish pLPAC genes from other

Notwithstanding the foregoing, in another embodiment, the pLPAC gene

species, and most preferred from a species in Brassicaceae (or Cruciferae).

This invention also provides isolated polypeptide products of the open reading frames of SEQ ID NO:1 or SEQ ID NO:10, having at least about 70% (preferably 80% and most preferably 90%) sequence identity, or at least about 80% similarity (preferably 90% and more preferably 95%) with the amino acid sequence of SEQ ID NO:2. In another embodiment, the polypeptides of the invention are at least about 40% identical (preferably 50%, and most preferably 60%) to the regions of residues 1-76, 613-669 or 1144-1161 of SEQ ID NO:2. Because of the natural sequence variation likely to exist among *plPAC* genes, one skilled in the art would expect to find up to about 40-40% nucleotide sequence variation, while still maintaining the unique properties of the *plPAC* gene and encoded polypeptide of the present invention. Such an expectation is due in part to the degeneracy of the genetic code, as well as to the known evolutionary success of conservative amino acid sequence variations, which do not appreciably alter the nature of the encoded protein. Accordingly, such variations are considered substantially the same as the invention and are included within the scope of the present invention.

Also provided are transgenic plants transformed with polynucleotides or nucleic acids of the invention.

Embodiments of the invention are described below, and in a most preferred

gene may be placed under a powerful constitutive promoter, such as the Cauliflower Mosaic Virus (CaMV) 35S promoter or the figwort mosaic virus 35S promoter. In a preferred embodiment, the 35S CaMV promoter is used. Transgenic plants expressing the *plPAC* gene under an inducible promoter (either its own promoter or a heterologous promoter) are also contemplated to be within the scope of the present invention. Inducible plant promoters include the tetracycline repressible/operator controlled promoter. In a preferred embodiment, a native *plPAC* promoter is used, and in a most preferred embodiment, residues 1-3429 of SEQ ID NO:10 is used. Plant species that are contemplated for overexpression of a *plPAC* coding sequence include, but are not limited to, soybean.

In another embodiment, overexpression of *plPAC* is induced to generate a co-suppression effect. This excess expression serves to promote down-regulation of both endogenous and exogenous *plPAC* genes.

In some instances, it may be desirable to down-regulate or inhibit expression of endogenous *plPAC* in plants possessing the gene. Accordingly, *plPAC* nucleic acid molecules, or fragments thereof, may also be utilized to control the production of *plPAC*-encoded P-glycoproteins. In one embodiment, full length *plPAC* antisense molecules or antisense oligonucleotides, targeted to specific regions of

preferred embodiments of the invention are provided in

upon transcription, produces the antisense sequences. Such constructs can be designed to produce full-length or partial antisense sequences. One example of antisense *plPAC* transgenic plants is given in Example 3.

5 In another embodiment, knock-out plants are obtained by screening a T-DNA mutagenized plant population for insertions in the *plPAC* gene (see Krysan et al., 1996, PNAS 93:145). One example of this embodiment of the invention is found in Example 3. Optionally, transgenic
10 plants can be created containing mutations in the region encoding the active site of *plPAC*. These last two embodiments are preferred over the use of anti-sense constructs due to the high homology among P-glycoproteins. The promoter of *ATPAB* is also provided in accordance with the
15 invention. This promoter has the useful properties of root expression and inducibility by NPPB. Presence of NPPB in the growth medium of *Arabidopsis* seedlings results in increased expression of *ATPAB* of the present invention.

Further, when approximately 4kb of upstream *ATPAC*
20 promoter DNA is fused to the *GUS* reporter gene and transformed into wild-type plants, *GUS* staining is strong in the hypocotyl of etiolated seedlings, but not in light grown seedlings. Further, expression is high in cotyledons, meristems, root, and the first true leaves of seedlings.
25 Staining was also observed in flowers and the apical portion

of the root. The expression of the *plPAC* gene in the plant species will

these promoter regions can easily be isolated from the *plPAC* genes that are provided with the invention, all plant *plPAC* gene promoters are provided with the invention. The nucleic acids of the invention therefore include a nucleic acid molecule that is at least about 70% identical (preferably 80% and most preferably 90%) to the residues 1-3429 of SEQ ID NO:10. Also provided are nucleic acids that hybridize to the nucleic acid residues 1-3429 of SEQ ID NO:10 preferably under moderate stringency, more preferably, high stringency, and most preferably, very high stringency).

Thus, the *plPAC* of the present invention encodes an ABC transporter that binds NPA and is involved with auxin transport in the plant. Mutants of *Arabidopsis* lacking *ATPAC* and double mutants lacking both *ATPAC* and *AtPGP1* display morphological phenotypes consistent with their demonstrated impairments in polar auxin transport. It has been widely accepted that NPA-sensitive regulatory site and the auxin-conducting channel of the efflux carrier are separate molecular entities. Strong evidence indicates that PIN-like genes encode the auxin-conducting channel of the efflux carrier (Galm et al., 1999, Curr. Op. Plant Biol. 2:375-381). However, as evidenced in Example 4 of the present invention, NPA-like genes are components of the NPA-sensitive regulatory site.

Expression of the *plPAC* gene of the present

invention in plants results in plants that are resistant to NPA. The strategy of the present invention is to provide a plant with a *plPAC* gene of the present invention and a promoter that drives expression of the *plPAC* gene in the plant.

induced by treatment of some dicots with herbicidal levels of the auxin analog, 2,4-D and in *Arabidopsis* by treatment with auxin transport inhibitors (Sieburth, (1999) *Plant Physiol.* 121: 1189-1193). This indicates that ATPAC of the present invention pumps auxin or auxin conjugates from sites of synthesis, such as the apical meristem and expanding cotyledons (Sachs, 1991 *Development* S1: 833-893). Under this model, tissues that express this pump would accumulate auxin as a result of the mutation and the altered auxin balance could be responsible for the altered growth patterns typifying the ATPAC phenotype. Further support of this model is the similarity of auxin to indolic substrates pumped by human MDR1, and the finding that ATPAC expression is increased by auxin. Also, the fact that NPA binds to ATPAC and that *atpac* knockout mutants can be phenocopied by auxin application suggests that ATPAC is an important component of the auxin transport and distribution machinery.

The present invention also provides antibodies capable of immunospecifically binding to polypeptides of the invention. In a preferred embodiment, the antibodies react immunospecifically with various epitopes of the *plPAC*-encoded polypeptides. In a particularly preferred embodiment, the antibodies are immunospecifically specific to the polypeptide of residues 1-76, 813-822 or 1144-1161 of SEQ ID NO:2.

The following examples sets forth the general principles of the invention.

EXAMPLE 1
Construction of the *plPAC* gene and expression of the *plPAC* protein in *Escherichia coli*

al., Molecular Cloning, Cold Spring Harbor Laboratory (1989) (hereinafter "Sambrook et al.") or Ausubel et al. (eds) Current Protocols in Molecular Biology, John Wiley & Sons (2001) (hereinafter "Ausubel et al.") are used.

5

**III. Preparation of PIPAC Nucleic Acid Molecules,
encoded Polypeptides, Antibodies Specific for the
Polypeptides and Transgenic Plants**

10

1. Nucleic Acid Molecules

15

PIPAC nucleic acid molecules of the invention may be prepared by two general methods: (1) they may be synthesized from appropriate nucleotide triphosphates, or (2) they may be isolated from biological sources. Both methods utilize protocols well known in the art.

20

The availability of nucleotide sequence information, such as the cDNA having SEQ ID NO:1, enables preparation of an isolated nucleic acid molecule of the invention by oligonucleotide synthesis. Synthetic oligonucleotides may be prepared by the phosphoramidite method employed in the Applied Biosystems 384 DNA Synthesizer or similar devices. The resultant construct may be purified according to methods known in the art, such as high performance liquid chromatography (HPLC). Long, double-stranded polynucleotides, such as a DNA molecule of the present invention, may be synthesized in stages, due to the

25

sequence of the nucleic acid molecule of the invention may be determined by sequencing the

appropriate cohesive termini for attachment of an adjacent segment. Adjacent segments may be ligated by annealing cohesive termini in the presence of DNA ligase to construct an entire long double-stranded molecule. A synthetic DNA molecule so constructed may then be cloned and amplified in an appropriate vector.

plPAC genes also may be isolated from appropriate biological sources using methods known in the art. In fact, the *ATPAC* clone was isolated from an *Arabidopsis* cDNA library using a partial clone obtained from *Brassica napus*. In alternative embodiments, genomic clones of *plPAC* may be isolated.

In accordance with the present invention, nucleic acids having the appropriate level sequence homology with part or all the coding regions of SEQ ID NO:1 or SEQ ID NO:10 may be identified by using hybridization and washing conditions of appropriate stringency. For example, hybridizations may be performed, according to the method of Sambrook et al., using a hybridization solution comprising:
5X SS_{0.1}, 1X Denhardt's reagent, 1.0% SDS, 100 µg/ml denatured, fragmented salmon sperm DNA, 0.05% sodium pyrophosphate and 0.01% formamide. Hybridization is carried out at 42°C for at least six hours. Following hybridization, filters are washed as follows: (1) 5 minutes at room temperature in 1X SS_{0.1} and 1% SDS; (2) 15 minutes at

room temperature in 0.1X SS_{0.1} and 1% SDS. The stringency conditions

molecules of specified sequence homology (Sambrook et al., 1989):

$$T_m = 81.5^{\circ}\text{C} - 16.6\log[\text{Na}^+] - 0.41(\%\text{G+C}) - 0.63(\%\text{formamide}) - 600/\text{\#bp in duplex}$$

5

As an illustration of the above formula, using $[\text{Na}^+] = [0.368]$ and 50% formamide, with GC content of 42% and an average probe size of 200 bases, the T_m is 57°C . The T_m of a DNA duplex decreases by $1 - 1.5^{\circ}\text{C}$ with every 1% decrease in homology. Thus, targets with greater than about 75% sequence identity would be observed using a hybridization temperature of 42°C .

10

The stringency of the hybridization and wash depend primarily on the salt concentration and temperature of the solutions. In general, to maximize the rate of annealing of the probe with its target, the hybridization is usually carried out at salt and temperature conditions that are 20-25 $^{\circ}\text{C}$ below the calculated T_m of the of the hybrid. Wash conditions should be as stringent as possible for the degree of identity of the probe to the target. In general, wash conditions should be approximately 12-20 $^{\circ}\text{C}$ below the T_m of the hybrid. As regards to the nucleic acids of the current invention, moderate stringency hybridization is defined as hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 0.1% heat-denatured salmon sperm DNA at 42°C ,

20

25

XXXXXX
XXXXXX

0.5% SDS at 65°C for 15 minutes. A very high stringency hybridization is defined as hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 µg/ml denatured salmon sperm DNA at 42°C, and wash in 0.1X SSC and 0.5% SDS at 65°C for 15 minutes.

Nucleic acids of the present invention may be maintained as DNA in any convenient cloning vector. In a preferred embodiment, clones are maintained in plasmid cloning-expression vector, such as pGEM-T (Promega Biotech, Madison, WI) or pBluescript (Stratagene, La Jolla, CA), either of which is propagated in a suitable *E. coli* host cell.

plPAC nucleic acid molecules of the invention include cDNA, genomic DNA, RNA, and fragments thereof which may be single- or double-stranded. Thus, this invention provides oligonucleotides (sense or antisense strands of DNA or RNA) having sequences capable of hybridizing with at least one sequence of a nucleic acid molecule of the present invention, such as selected segments of SEQ ID NO:1 or SEQ ID NO:10. Such oligonucleotides are useful as probes for detecting plPAC mRNA in test samples, e.g. by PCR amplification, to screen for genes or for the positive or negative regulation of expression of plPAC genes at or before translation of the mRNA into proteins.

The plPAC vector is also expected to be useful in

transforming cells with plPAC. The plPAC vector may be used to transform cells with plPAC or to transform cells with plPAC and a reporter gene.

used in chimeric gene constructs to facilitate inducible expression of any coding sequence of interest, upon exposure to NPPB or similar acting compounds.

5

2. Proteins and Antibodies

Polypeptides encoded by *plPAC* nucleic acids of the invention may be prepared in a variety of ways, according to known methods. If produced *in situ* the polypeptides may be purified from appropriate sources, e.g., plant roots or other
10 plant parts.

Alternatively, the availability of nucleic acid molecules encoding the polypeptides enables production of the proteins using *in vitro* expression methods known in the art. For example, a cDNA or gene may be cloned into an appropriate
15 *in vitro* transcription vector, such as pSP64 or pSP65 for *in vitro* transcription, followed by cell-free translation in a suitable cell-free translation system, such as wheat germ or rabbit reticulocytes. *In vitro* transcription and translation systems are commercially available, e.g., from Promega
20 Biotech, Madison, Wisconsin or BRL, Rockville, Maryland. According to a preferred embodiment, larger quantities of *plPAC*-encoded proteins may be produced by expression in a suitable prokaryotic or eukaryotic system. For example, part or all of a DNA molecule, such as the cDNA having SEQ ID
25 NO:1, may be inserted into a plasmid vector adapted for expression in a prokaryotic or eukaryotic system.

For the purpose of the present invention, the methods necessary for expression of the proteins of the invention are well known in the art.

manner as to permit expression of the DNA in the host cell. Such regulatory elements required for expression include promoter sequences, transcription initiation sequences and, optionally, enhancer sequences.

5 The pLPAC polypeptide produced by gene expression in a recombinant prokaryotic or eucaryotic system may be purified according to methods known in the art. In a preferred embodiment, a commercially available expression/secretion system can be used, whereby the
10 recombinant protein is expressed and thereafter secreted from the host cell, to be easily purified from the surrounding medium. If expression/secretion vectors are not used, an alternative approach involves purifying the recombinant protein by affinity separation, such as by immunological
15 interaction with antibodies that bind specifically to the recombinant protein. Such methods are commonly used by skilled practitioners.

 The pLPAC-encoded polypeptides of the invention, prepared by the aforementioned methods, may be analyzed
20 according to standard procedures.

 Polypeptides and clonal antibodies directed toward any of the polypeptides encoded by pLPAC may be prepared according to standard methods. Monoclonal antibodies may be prepared according to general methods of Köhler and Milstein,
25 following standard protocols.

 The polypeptides and antibodies of the invention may be used in a variety of assays. These include, but are not limited to, immunoassays, such as enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and Western blots.

limited to, *Agrobacterium* vectors, PEG treatment of protoplasts, biolistic DNA delivery, UV laser microbeam, gemini virus vectors, calcium phosphate treatment of protoplasts, electroporation of isolated protoplasts, agitation of cell suspensions with microbeads coated with the transforming DNA, direct DNA uptake, liposome-mediated DNA uptake, and the like. Such methods have been published in the art. See, e.g., Methods for Plant Molecular Biology (Weissbach & Weissbach, eds., 1988); Methods in Plant Molecular Biology (Schuler & Zielinski, eds., 1989); Plant Molecular Biology Manual (Selvin, Schilperoort, Verma, eds., 1993); and Methods in Plant Molecular Biology - A Laboratory Manual (Maliga, Klüssig, Cashmore, Gruissem & Varner, eds., 1994).

The method of transformation depends upon the plant to be transformed. The biolistic DNA delivery method is useful for nuclear transformation. In another embodiment of the invention, *Agrobacterium* vectors are used to advantage for efficient transformation of plant nuclei.

In a preferred embodiment, the gene is introduced into plant nuclei using *Agrobacterium* binary vectors. Such vectors include, but are not limited to, BIN19 (Beyan, 1984, Nucleic Acids Res. 12:4721) and derivatives thereof, the pBI vector system (Chilton et al., 1987, PNAS 83:844751), and binary vectors such as Ma-1 and pGA492 (An, 1986) and others (for a review, see e.g., Methods Mol Biol 44:17-28).

The gene is operably linked to a promoter and a polyA signal sequence. The gene is operably linked to a promoter and a

(e.g., promoters and translational regulatory sequences) and 3' regulatory sequences (e.g., terminators).

DNA constructs for transforming a selected plant comprise a coding sequence of interest operably linked to appropriate 5' and 3' regulatory sequences (e.g., promoters and translational regulatory sequences) and 3' regulatory sequences (e.g., terminators). In a preferred embodiment, the coding region is placed under a powerful constitutive promoter, such as the Cauliflower Mosaic Virus (CaMV) 35S promoter or the figwort mosaic virus 35S promoter. Other constitutive promoters contemplated for use in the present invention include, but are not limited to: T-DNA mannopine synthetase, nopaline synthase (NOS) and octopine synthase (OCS) promoters.

Transgenic plants expressing a sense or antisense SDS coding sequence under an inducible promoter are also contemplated to be within the scope of the present invention. Inducible plant promoters include the tetracycline repressor/operator-controlled promoter, the heat shock gene promoters, stress (e.g., wounding)-induced promoters, defense responsive gene promoters (e.g. phenylalanine ammonia lyase genes), wound-inducible promoters (e.g. hydroxyproline rich cell wall glycoprotein genes), chemically-inducible gene promoters (e.g. chitinase genes, glucanase genes, chitinase genes, etc.) and dark-inducible gene promoters (e.g., asparagine synthetase gene) to name a few.

The present invention and development-specific promoters are

also contemplated to be within the scope of the present invention. Such promoters include, but are not limited to, promoters for the following proteins: SDS, etc.

promoters for expression in photosynthetic tissue; the various seed storage protein gene promoters for expression in seeds; and the tissue-specific glutamine synthetase gene promoters where expression in roots is desired.

5 The coding region is also operably linked to an appropriate 3' regulatory sequence. In a preferred embodiment, the glutamine synthetase polyadenylation region (NOS) is used. Other useful 3' regulatory regions include, but are not limited to the octopine (OCS) polyadenylation
10 region.

 Using an *Agrobacterium* binary vector system for transformation, the *pLFA3* coding region, under control of a constitutive or inducible promoter as described above, is linked to a nuclear drug resistance marker, such as kanamycin
15 resistance. *Agrobacterium*-mediated transformation of plant nuclei is accomplished according to the following procedure:

 (1) the gene is inserted into the selected *Agrobacterium* binary vector;

 (2) transformation is accomplished by co-cultivation of plant tissue (e.g., leaf discs) with a
20 suspension of the selected *Agrobacterium*, followed by incubation of the explants on growth medium in the absence of the drug marker in the selective medium (see, e.g., Horsch et al. 1988, *Chloroplast Harb Symp Quant Biol.* 50:4337);

25 (3) the explant is then transferred onto the selective medium to identify transformed tissue; and

 (4) the transformed tissue is then grown to maturity and the amount of protein produced is determined. The tissue specificity of expression is

of the *plPAC* gene in transformed plants can vary depending on the position of the insertion into the nuclear genome. Such position effects are well known in the art. For this reason, several nuclear transformants should be regenerated and tested for expression of the transgene.

IV. Uses of *plPAC* Nucleic Acids, Encoded Proteins and Antibodies

1. *plPAC* Nucleic Acids

plPAC nucleic acids may be used for a variety of purposes in accordance with the present invention. The DNA, RNA, or fragments thereof may be used as probes to detect the presence or absence of expression of *plPAC* genes. Methods in which *plPAC* nucleic acids may be utilized as probes for such assays include, but are not limited to: (1) *in situ* hybridization; (2) Southern hybridization (3) northern hybridization; and (4) assorted amplification reactions such as polymerase chain reactions (PCR).

The *plPAC* nucleic acids of the invention may also be utilized as probes to identify related genes from other plant species. As is well known in the art and described above, hybridization conditions may be adjusted to allow hybridization between a probe and probes with complementary sequences of varying degrees of homology. Thus, *plPAC* nucleic acids may be used to advantage to identify and characterize genes of varying degrees of relationship to the *plPAC* gene.

Very respectfully,

Attorney for Applicant

5

2. ELPAC Proteins and Antibodies

10

20

and purifying such proteins. For example, antibodies may be utilized for activity separation of proteins with which they immunospecifically interact. Antibodies may also be used to immunoprecipitate proteins from a sample containing a mixture of proteins and other biological molecules.

3. plPAC Transgenic Plants

Transgenic plants that over- or under- express plPAC can be used in a variety of agronomic and research applications. From the foregoing discussion, it can be seen that plPAC and the homologs, and transgenic plants containing them will be useful for improving stress resistance or tolerance in plants. This provides an avenue for developing marginal or toxic soil environments for crop production. Both over- and under-expressing plPAC transgenic plants have great utility in the research of herbicides and other xenobiotic compounds.

As discussed above and in greater detail in Example 1, the similarity between plant and mammalian *mdr* genes indicates that their functional aspects will also be conserved. It is thus expected to play an important role in the export of toxic metabolic or xenobiotic compounds from cells. The fact that plPAC also is inducible and appears to be constitutively expressed in roots, where contact with such compounds often occurs, makes plPAC particularly useful in the genetic engineering of plants to

apply to the development of plants having increased resistance to

of the kinds of compounds that should be detoxified by the *plPAC*. The inhibitors include, but are not limited to, hydrophobic (i.e., lipophilic) herbicides and other compounds, such as 3,4-dichlorophenyl-1,1-dimethyl urea (also known as DCMU oruron, available from Sigma Chemical Co., St. Louis, Mo.) or other hydrophobic compounds that disrupt photosynthetic electron transport, as well as Metachlor (Ciba Geigy, Basel Switzerland), Taurocholate (Sigma Chemical Co.), Primisulfuron (Ciba Geigy), and IRL-1803.

As illustrated in Example 2, plant cells that over-express a *plPAC* gene have surprisingly higher growth rate with or without the xenobiotic compound Rhodamine 6G. It is contemplated that *plPAC* overexpression may be a generally useful way to increase plant and plant cell culture growth, even without the presence of xenobiotic compounds.

In addition to the above-mentioned features and advantages of transgenic plants that are altered in their expression of *plPAC*, these plants will also be altered in auxin transport. Through the use of developmental or tissue specific promoters, plants having a pre-determined alteration in auxin transport may be produced, providing agronomically or horticulturally desired traits/features to such plants.

The following specific examples are provided to illustrate embodiments of the invention. They are not intended to limit the scope of the invention, which is defined by the claims.

EXAMPLE 1

Isolation and Analysis of a *plPAC* Gene From *Arabidopsis thaliana*

The *ATPAC* of the present invention was identified by its up-regulation in response to a chloride ion channel blocker. *Brassica napus* plants were grown either in the presence or absence of 2 μ M 5-nitro-2-(3-phenylpropylamino) benzoic acid (NPPB). After five days, the roots of the seedlings were harvested and total RNA was extracted separately from the treated and untreated plants. From the total RNA preparations, poly (A)⁺ RNA was isolated and used as the starting material to create a cDNA subtraction library, using the CLONTECH PCR-SELECT™ cDNA Subtraction Kit and accompanying instructions (CLONTECH Laboratories, Inc., Palo Alto, CA).

Using the subtractive hybridization kit, a gene fragment was identified that was up-regulated in response to treatment of the plants with NPPB. This fragment (0.5 kb) was used to screen a cDNA library of *Arabidopsis thaliana*, from which a full-length cDNA clone was isolated. The nucleotide sequence of this cDNA clone, referred to as ATPAC (*Arabidopsis thaliana* putative anion channel) is set forth below as SEQ ID NO:1.

The *atpac* cDNA clone encodes a polypeptide 1,254 amino acids in length. The deduced amino acid sequence encoded by SEQ ID NO:1 is shown in Figure 1 as "atpac" (SEQ ID NO:1), in a library with the following sequences: (1) hmdr1 (SEQ ID NO:3); (2) hmdr2 (SEQ ID NO:4); (3) hmdr3 (SEQ ID NO:5); (4) hmdr4 (SEQ ID NO:6); (5) atpac1 (SEQ ID NO:7); and

that *Arabidopsis thaliana* has a family of anion channel proteins that *Arabidopsis thaliana* has a family of the members of

ABC transporters. In none of the databases, including the EST collection, does an exact match exist. The ABC transporter family is very large, consisting of at least two sub-groups, *mrp* and homologs and *mdr* and homologs. The only examples of plant *mdr*-like genes are *atpgp1* and *atpgp2* from *A. thaliana* and two homologs from potato and barley, respectively. Though the *atpgp1* and *atpgp2* genes are similar to ATPAC, they are only 41 and 50% identical, respectively, indicating that ATPAC is a distinct gene by comparison.

Sequence homology with the potato and barley *mdr*-like genes is even more divergent. Another difference between the *atpgp1* gene and the ATPAC gene is their respective preferential expression in inflorescens and roots, respectively.

EXAMPLE 2

Effect of ATPAC Expression in Bacterial Cells on Their Ability to Detoxify Rhodamine 6G

The compound Rhodamine 6G is a well known substrate of mammalian p-glycoproteins (Kolaczowski et al., J. Biol. Chem., 263:3194-3197, 1988). The ability of a cell to detoxify the compound is indicative of activity of p-glycoproteins. A bacterial cell line was transformed with an expression vector comprising ATPAC. The growth rate of transformed and non-transformed cells was then measured, in the presence or absence of Rhodamine 6G. Results are shown

in Table 1. The growth rate of the transformed cells was significantly higher than that of the non-transformed cells in the presence or absence of

Rhodamine 6G. These results demonstrate that ATPAC encodes a functional and robust p-glycoprotein.

Example 3
Transgenic Plants that Overexpress
and Underexpress ATPAC

Transformation construct. The *Agrobacterium* binary vector pP211 (Hajdukiewicz et al., 1994 Plant Mol. Biol. 25:989-994) was digested with *EcoRI* and *SmaI*, and selfligated. This molecule was named pP211'. The *Agrobacterium* binary vector pGEN7366 (Clontech, CA) was digested with *XhoI* and cloned in *SalI* digested pP211'. We named this binary vector pPZPPCGN. The 3.2 kb full-length ATPAC cDNA was cloned into the pGHI2 vector. After digestion with *SmaI* (in the multiple cloning site upstream) and *EcoRI*, a 3.1 kb cDNA fragment was cut out. This *SmaI/EcoRI* 3.1 kb fragment was cloned into the *SmaI/EcoRI* site of pPZPPCGN. The rest of ATPAC gene was amplified using polymerase chain reaction to have translationally fused HA tag at its 3' terminal. After ligating *EcoRI* linkers to the ends of the resulting PCR product, the 0.7 kb fragment was cloned into the *EcoRI* site of the *SmaI/EcoRI* 3.1 kb cDNA fragment in pPZPPCGN. The final construct was named pATPACOE.

Plant transformation. pATPACOE was introduced into *Agrobacterium tumefaciens* strain by a direct transformation method. *Agrobacterium* mediated transformation was performed using the method of Clontech (Clontech et al., 1997, GE Acad.

T2 seeds. T3 seeds were selected from kanamycinresistant T2 plants. T3 plants which showed 100% kanamycinresistance were selected and were considered homozygous for the transgene.

Antisense Plants. The full length cDNA in pBluescript SK(+) vector (Stratagene, CA) is digested with *EcoRI* (there is a cleavage site in the upstream polylinker) and *SspI*. The resulting 1.3 Kb fragment representing a 5' portion of the *AtPAC* cDNA was cloned into the aforementioned pPZP001, which is then digested with *EcoRI/SmaI*, ensuring that this fragment of the cDNA was inserted in the antisense orientation. This construct was named pATPACAE. pATPACAE was introduced into *Arabidopsis* plants by *Agrobacterium* transformation, as described above.

Knock-out Plants. The method of Krysan et al (1996, *PNAS* 93:8148, incorporated by reference herein) was followed using the following primers:

Genespecific primers:

AtpacF: TACTGCTCAGTATCTCTTTTCTCACTA (SEQ ID NO:11)

AtpacR: TTGAATCATAAATCAATCAACACCTC (SEQ ID NO:12)

Primers for TDNA left border:

JL202: TTTTATAATTTTATTTTACATTTAC (SEQ ID NO:13)

JL270: TTTTATAATTTTATTTTACATTTAC (SEQ ID NO:14)

T-DNA insertion mutants were isolated by PCR-based screen of DNA from 12 alleles of ATPAC and one ATPGP1 allele was isolated. At the seedling stage, both alleles of ATPAC were present in the embryonic cotyledons and

were somewhat curved and wrinkled along the margin. Bolting of the inflorescence stem was delayed by 2.8 days on average, relative to wild type. The bolt grew more slowly than wild type and started to die when the wild type length was

5 ultimately reached. These phenotypes coincided with the sites of expression indicated by GUS staining (as described herein). None of these phenotypes were present in plants transformed with a minimal fragment containing the wild type *ATPAC* promoter and coding sequence ("*atpac1-1*"). This mutant
10 did not display any overt phenotype as a seedling or an adult plant. Double mutants were constructed by crossing *atpac1-1* and *atpg1-1* plants. F1 individuals appeared wild type and were permitted to self pollinate. Approximately one in sixteen of the F2 seedlings displayed extremely down-curved
15 cotyledons when grown in the light and also displayed shorter, wavy hypocotyls when grown in the dark. PCR analysis confirmed that these seedlings were homozygous double mutants. In adult double mutant plants were severely stunted in growth. Primary inflorescence stems of the double
20 mutants and floral pedicels were also wavy in appearance, indicating that the direction of growth periodically changed during development of these organs. After 12 hours of growth the double mutant produced abundant secondary inflorescence stems, indicating a large reduction in apical
25 dominance. Fertilization of flowers in the double mutants was possible due to the elongation of the stamen filament.

Example 4

Effect of Auxin (IAA) on ATPAC Expression

Expression in Yeast and Xenopus Oocytes. ATPAC

5 cDNA was expressed in wild type yeast as well as in yeasts lacking even ABC transporters (as described by Decottignies et al., 1993) (J. Biol. Chem. 273:12612-22) in order to create a heterologous system for studying function of the transporter. The plants were exposed to toxic compounds that
10 are known substrates for human MDR1. ATPAC did not confer any measurable resistance to the toxic substrates. Further, there was no evidence of a drug-pumping role for ATPAC.

In order to examine whether ATPAC functions as an anion channel or a regulator of an anion channel,
15 complementary RNA was from an ATPAC cDNA template was injected into Xenopus oocytes to produce a heterologous expression system amenable to electrophysiology. No currents associated with ATPAC were observed by two electrode voltage clamping.

20 Treatment of wild type seedlings with 2,4-D or high concentrations of auxin resulted in plants having strongly curved stems.

ATPAC binds the Auxin Transport Inhibitor naphthylphthalamic acid (NPA). Yeast expressing ATPAC was
25 assayed for NPA binding. NPA bound tightly and specifically to ATPAC expressed in yeast but not to control yeast. Bound NPA was released by treatment with a related but non-binding

30 compound, 2,4-dichlorophenoxyacetic acid, indicating that ATPAC is a specific auxin transport inhibitor.

NPA binds to ATPAC and that *atpac* knockout mutants can be phenocopied by auxin application suggests that ATPAC is an important component of the auxin transport and distribution machinery.

5 Effect of ATPAC and AtPGP1 on Auxin Transport.

Three different auxin transport assays of the *Ws* wild type, ATPAC-1, ATPAC-2 and *atpgp1* were performed. The first measured the basipetal movement of auxin in seedlings as described by Murphy et al. (2000) *Planta* 211:315-324. A 0.1
10 μl microdroplet of radioactive auxin was placed on the apex of a light-grown seedling. Four hours later, the amount of radioactivity collected on moist filter paper that contacted the root tip was determined. Polar auxin transport measured in this manner was severely reduced in both alleles of ATPAC
15 and especially in the double mutant, but not in the *atpgp1* mutant.

The second assay measured the basipetal transport of radioactive auxin through etiolated seedlings inverted in a reservoir containing radioactive auxin (Garbers et al.
20 (1996) *PLoS J.* 15:2116-2119). The results were similar to those with the first assay described above.

The third assay measured basipetal transport of auxin in the inflorescence stem in a method described by Ruegger et al. (1999) *Plant Cell* 9:745-57. A segment of
25 inflorescence stem was excised and immersed apical-end down in a tube containing a known volume of radioactive auxin. At
30 the time of excision, the inflorescence was exposed to a 10 min pulse of auxin.

40 Auxin transport in the inflorescence stem of *atpgp1* and *atpac*

significantly increased transport in tissue segments taken from the lower portion of the inflorescence. This is indicative of gradients of function of at least two MDR-like gene products along the inflorescence axis.

5

While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be limited to such embodiments. Various modifications may be

10

made thereto without departing from the scope and spirit of the present invention, as set forth in the following claims.